

NO-A188 495

INSTRUMENTATION FOR COLLISIONAL ENERGY TRANSFER STUDIES

1/1

(U) WISCONSIN UNIV-MADISON DEPT OF CHEMISTRY F F CRIM


27 OCT 87 AFOSR-TR-87-1786 AFOSR-86-0244

UNCLASSIFIED

F/G 20/5

ML





RESOLUTION TEST CHART

REPORT DOCUMENTATION PAGE

AD-A188 495

DTIC

10. RESTRICTIVE MARKINGS

20. DECLASSIFICATION/DOWNGRADING SCHEDULE
N/A

DEC 1 1987

3. DISTRIBUTION AVAILABILITY OF REPORT
Approved for public release;
Distribution unlimited

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

D

5. MONITORING ORGANIZATION REPORT NUMBER(S)
AFOSR-TR- 87-1706

6a. NAME OF PERFORMING ORGANIZATION

University of Wisconsin

6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

AFOSR/NC

6c. ADDRESS (City, State and ZIP Code)

Department of Chemistry
1101 University Avenue
Madison, WI 53706

7b. ADDRESS (City, State and ZIP Code)

Bldg. 410
Bolling AFB, D. C. 20332

8a. NAME OF FUNDING SPONSORING ORGANIZATION

AFOSR

8b. OFFICE SYMBOL
(If applicable)

NC

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

AFOSR-86-0244

8c. ADDRESS (City, State and ZIP Code)

Bldg. 410
Bolling AFB 20332-6448

10. SOURCE OF FUNDING NOS

PROGRAM
ELEMENT NOPROJECT
NO.TASK
NO.WORK UNIT
NO.

61102F

2319

A2

11. TITLE (Include Security Classification) Instrumentation for
Collisional Energy Transfer Studies

12. PERSONAL AUTHOR(S)

F. F. Crim

13a. TYPE OF REPORT

Final

13b. TIME COVERED

FROM 7/15/85 TO 7/14/87

14. DATE OF REPORT (Yr., Mo., Day)

1987/10/27

15. PAGE COUNT

3

16. SUPPLEMENTARY NOTATION

17. COSAT CODES

FIELD

GROUP

SUB. GR.

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Molecular energy transfer, Vibrational energy transfer

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

Two dye laser systems were purchased for use in energy transfer studies and in unimolecular decomposition experiments. Individual angular momentum states of acetylene in the fourth vibrational state of the CH stretching mode were prepared. It was found that these molecules which contain 10,000 wave numbers of vibrational energy relax at several times the gas kinetic collision rate. There is a striking dependence on the initial rotational state excited implying relaxation of only relatively small amounts of rotational energy.

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☒ DTIC USERS ☐

21. ABSTRACT SECURITY CLASSIFICATION

Unclassified

22a. NAME OF RESPONSIBLE INDIVIDUAL

F. J. Wodarczyk

22b. TELEPHONE NUMBER

(202) 767-4960

22c. OFFICE SYMBOL

NC

Final Report

Grant No. AFOSR-86-0244

(15 July 1986 to 14 July 1987)

This grant is part of the University Research Instrumentation Program (URIP) that allowed us to advance our DoD sponsored research very significantly by purchasing lasers for use in our collisional energy transfer studies and in our unimolecular decomposition experiments. These benefits have come not only from the use of the hardware on our AFOSR sponsored projects but also by reducing competition for equipment by other projects. The net result has been some dramatic new collisional energy transfer results and spectroscopic measurements in our AFOSR sponsored project. In other DoD sponsored research, we have obtained new data in two photodissociation experiments and a new vibrational overtone spectroscopic measurement.

The primary goal of our URIP proposal was to advance our studies of collisional energy transfer in highly vibrationally excited polyatomic molecules. We have now succeeded in preparing individual angular momentum states of acetylene (C_2H_2) in the region of the third overtone of the CH stretching transition $3\nu_{CH}$ and monitoring the resulting highly vibrationally excited molecule using ultraviolet laser induced fluorescence. The experiments have three primary aspects: they explore the nature of the highly vibrationally excited state produced by vibrational overtone excitation, they exploit the two-photon combination of vibrational excitation and laser induced fluorescence to excite directly

previously unobservable vibrations in the electronically excited molecule, and they follow the collisional relaxation of individual quantum states in highly vibrationally excited molecules. The collisional energy transfer studies have shown that these molecules that contain $10\,000\text{ cm}^{-1}$ of vibrational energy relax at several times the gas kinetic collision rate. We note a striking dependence in the probe excitation spectra on the initial rotational state excited and have tentatively concluded that the relaxation proceeds by extremely rapid vibrational energy transfer with only relatively small amounts of rotational relaxation. We have carefully determined the rate of loss of the excited molecules and are now identifying the final states of the relaxation. We expect to sort out the mechanism of the collisional energy transfer and extend the work to other vibrational levels and molecules.

The other DoD sponsored research that has benefitted from the increased experimental time available because of the URIP hardware is a unimolecular decomposition experiment funded by the Army Research Office. We have now studied the vibrationally mediated photodissociation of HONO_2 and made the first state resolved measurements on its vibrational overtone induced decomposition. In the vibrationally mediated decomposition experiments, one photon excites an overtone vibration and another fragments the resulting vibrationally excited molecule by exciting it to a repulsive electronic surface. This approach has revealed unique photodissociation dynamics, as seen in the quantum state distribution of the products. We have also made the first state

resolved unimolecular decomposition measurements on this molecule by exciting the fourth ($5\nu_{\text{OH}}$) or fifth ($6\nu_{\text{OH}}$) OH stretching overtone vibration and probing the individual quantum states of the OH fragment.

APPROVED FOR
NPD GROUP
ITC TAB
UNIT 1000
SHOULD
By
Date
Doc
A-1

END

DATE

FILMD

3-88

DTIC